

beta-globin gene transfer to human bone marrow for sickle cell disease.

Journal: J Clin Invest

Publication Year: 2013

Authors: Zulema Romero, Fabrizia Urbinati, Sabine Geiger, Aaron R Cooper, Jennifer Wherley, Michael L Kaufman, Roger P Hollis, Rafael Ruiz de Assin, Shantha Senadheera, Arineh Sahagian, Xiangyang Jin, Alyse Gellis, Xiaoyan Wang, David Gjertson, Satiro Deoliveira, Pamela Kempert, Sally Shupien, Hisham Abdel-Azim, Mark C Walters, Herbert J Meiselman, Rosalinda B Wenby, Theresa Gruber, Victor Marder, Thomas D Coates, Donald B Kohn

PubMed link: 23863630

Funding Grants: Stem Cell Gene Therapy for Sickle Cell Disease

Public Summary:

Autologous hematopoietic stem cell gene therapy is an approach to treating sickle cell disease (SCD) patients that may result in lower morbidity than allogeneic transplantation. We examined the potential of a lentiviral vector (LV) (CCL-betaAS3-FB) encoding a human hemoglobin (HBB) gene engineered to impede sickle hemoglobin polymerization (HBBAS3) to transduce human BM CD34⁺ cells from SCD donors and prevent sickling of red blood cells produced by in vitro differentiation. The CCL-betaAS3-FB LV transduced BM CD34⁺ cells from either healthy or SCD donors at similar levels, based on quantitative PCR and colony-forming unit progenitor analysis. Consistent expression of HBBAS3 mRNA and HbAS3 protein compromised a fourth of the total beta-globin-like transcripts and hemoglobin (Hb) tetramers. Upon deoxygenation, a lower percentage of HBBAS3-transduced red blood cells exhibited sickling compared with mock-transduced cells from sickle donors. Transduced BM CD34⁺ cells were transplanted into immunodeficient mice, and the human cells recovered after 2-3 months were cultured for erythroid differentiation, which showed levels of HBBAS3 mRNA similar to those seen in the CD34⁺ cells that were directly differentiated in vitro. These results demonstrate that the CCL-betaAS3-FB LV is capable of efficient transfer and consistent expression of an effective anti-sickling beta-globin gene in human SCD BM CD34⁺ progenitor cells, improving physiologic parameters of the resulting red blood cells.

Scientific Abstract:

Autologous hematopoietic stem cell gene therapy is an approach to treating sickle cell disease (SCD) patients that may result in lower morbidity than allogeneic transplantation. We examined the potential of a lentiviral vector (LV) (CCL-betaAS3-FB) encoding a human hemoglobin (HBB) gene engineered to impede sickle hemoglobin polymerization (HBBAS3) to transduce human BM CD34⁺ cells from SCD donors and prevent sickling of red blood cells produced by in vitro differentiation. The CCL-betaAS3-FB LV transduced BM CD34⁺ cells from either healthy or SCD donors at similar levels, based on quantitative PCR and colony-forming unit progenitor analysis. Consistent expression of HBBAS3 mRNA and HbAS3 protein compromised a fourth of the total beta-globin-like transcripts and hemoglobin (Hb) tetramers. Upon deoxygenation, a lower percentage of HBBAS3-transduced red blood cells exhibited sickling compared with mock-transduced cells from sickle donors. Transduced BM CD34⁺ cells were transplanted into immunodeficient mice, and the human cells recovered after 2-3 months were cultured for erythroid differentiation, which showed levels of HBBAS3 mRNA similar to those seen in the CD34⁺ cells that were directly differentiated in vitro. These results demonstrate that the CCL-betaAS3-FB LV is capable of efficient transfer and consistent expression of an effective anti-sickling beta-globin gene in human SCD BM CD34⁺ progenitor cells, improving physiologic parameters of the resulting red blood cells.

Source URL: <http://www.cirm.ca.gov/about-cirm/publications/beta-globin-gene-transfer-human-bone-marrow-sickle-cell-disease>